Kropina spaces of constant curvature II.
(long version)

By
R. Yoshikawa and K. Okubo

arXiv:1110.5128v1 [math.DG] 24 Oct 2011

Introduction. In 1991, Matsumoto considered the necessary and sufficient conditions for a Kropina space to be of constant curvature and obtained

Theorem M([3], [4]) *An n-dimensional Kropina space is of constant curvature K, if and only if, putting $r_{ij} = (b_{i;j} + b_{j;i})/2$ and $s_{ij} = (b_{i;j} - b_{j;i})/2$, we have*
(1) $r_{ij} = Mb^2 a_{ij}$,
(2) $U_i = u_i + Mb_i$ *is a gradient vector field,*
(3) $w_{ij} = s_{ij}s^r{}_j$ *is written in the form*
$w_{ij} = b_i w_j + b_j w_i + (R - 2M^2b^2)b_ib_j - b^2U_iU_j - 4Ka_{ij}$,
(4) $s_{ij;k}$ *is written in the form*
$s_{ij;k} = A_{(ij)}\{a_{ik}[W_j + (R - M^2b^2)b_j] + b_iU_{jk}\}$,
where the symbol $A_{(ij)}$ denotes interchanges of indices i and j and subtraction,
(5) the curvature tensor R_{hijk} of the associated Riemannian space has the form
${}^{\alpha}R_{hikj} = A_{(kj)}\{Ra_{hk}a_{ij} - a_{hk}U_{ij} - a_{ij}U_{hk}\}$,
where we put $M = a^{ij}r_{ij}/nb^2$, $u_i = b_rs^r{}_i/b^2$, $U_{ij} = U_{i;j} + U_iU_j$, $w_i = u_rs^r{}_i$, $W_i = w_i + Mb^2U_i$ and $R = 4K/b^2 + U_rU^r$.

In [7], we characterized a Kropina space by some Riemannian space (M, h) and a vector field W of constant length 1 on it, and rewrite Theorem M using h and W. But, a few years after we noticed that we should replace (3) in Theorem M with the condition

$$w_{ij} = b_iw_j + b_jw_i + (R - 2M^2b^2)b_ib_j - b^2U_iU_j - 4Ka_{ij} + Mb^2(U_ib_j + U_jb_i)$$

which is equivalent to the equality in Lemma 2 of [3]. Then, it follows that the condition 4 of Theorem 2 in [7] should be omitted. Therefore, we consider the same problem in another way.

In this paper, we rewrite the condition for a Kropina space to be of constant curvature with a Riemannian metric h and a vector filed W and obtain the necessary and sufficient conditions for a Kropina space to be of constant curvature.

In the first section, we will show again that a Kropina space is characterized by some Riemannian metric h and a vector field W of constant length 1 on it, and in the second section, we will rewrite the coefficients of the geodesic spray in a Kropina space by the Riemannian metric h and the vector field W.

In Finsler geometry, the necessary and sufficient condition for a Kropina space to be of scalar curvature is well-known ([2]). We rewrite the condition by h and W, and using the relation of h and W we obtain the necessary and sufficient conditions for a Kropina space to be of constant curvature by straightforward calculations. Our main result is Theorem 4. This Theorem 4 is the corrected version of Theorem 2 in [7]. Therefore, this paper is the corrected version of [7].

1 The characterization of a Kropina metric.

Let (M, α) be an $n(\geq 2)$-dimensional differential manifold endowed with a Riemannian metric α. A Kropina space $(M, \alpha^2/\beta)$ is a Finsler space whose fundamental function is given by $F = \alpha^2/\beta$, where $\alpha = \sqrt{a_{ij}(x)y^iy^j}$ and $\beta = b_i(x)y^i$. Even though Kropina spaces can be studied in more general case ([4]), in this paper, we suppose that the matrix (a_{ij}) is positive definite.

Let us remark that for a Kropina space $(M, \alpha^2/\beta)$ the Kropina metric $F = \alpha^2/\beta$ can be rewrited as follows:

$$\frac{\alpha^2}{F^2} - \frac{\beta}{F} + \frac{b^2}{4} \quad = \quad \frac{b^2}{4}.$$

where $b^2 = a^{ij}b_ib_j$ and $(a^{ij}) = (a_{ij})^{-1}$. Let $\kappa(x)$ be a function of (x^i) alone. Multiplying the above equation by $e^{\kappa(x)}$, we have

$$(1.1) \qquad e^{\kappa(x)}a_{ij}\frac{y^i}{F}\frac{y^j}{F} - e^{\kappa(x)}a_{ij}\frac{y^i}{F}b^j + \frac{1}{4}e^{\kappa(x)}a_{ij}b^ib^j = \frac{e^{\kappa(x)}b^2}{4},$$

Defining a new Riemannian metric $h = \sqrt{h_{ij}(x)y^iy^j}$ and a vector field $W = W^i(\partial/\partial x^i)$ on M by

$$(1.2) \qquad h_{ij} = e^{\kappa(x)}a_{ij} \;\; \textit{and} \;\; 2W_i = e^{\kappa(x)}b_i,$$

where $W_i = h_{ij}W^j$, the equation (1.1) reduces to

$$\left|\frac{y}{F} - W\right| = |W|.$$

In the above equation, the symbol $|\cdot|$ denotes the length of a vector with respect to the Riemannian metric h.

We notice that the equation $|W| = 1$ holds if and only if the function $\kappa(x)$ satisfies the condition

$$(1.3) \qquad e^{\kappa(x)}b^2 = 4.$$

Suppose that the function $\kappa(x)$ satisfies (1.3), then we have $|W| = 1$ and

$$(1.4) \qquad \left|\frac{y}{F} - W\right| = 1.$$

Therefore, in each tangent space T_xM, the indicatrix of the Kropina metric necessarily goes through the origin.

Conversely, consider a Riemannian space (M, h), where $h = \sqrt{h_{ij}(x)y^iy^j}$, and a unit vector field $W = W^i(\partial/\partial x^i)$ on it. Then, we consider the metric F characterized by (1.4). Solving for F from (1.4), we get

$$F = \frac{|y|^2}{\{\sqrt{2}h(y, W)\}^2}.$$

Comparing the above equality with a Kropina metric $F = \alpha^2/\beta$, we obtain (1.2) and from the assumption $|W| = 1$ we get (1.3).

Summarizing the above discussion, we obtain

Theorem1 *Let (M, α) be an $n(\geq 2)$-dimensional Riemannian space with a Riemannian metric $\alpha = \sqrt{a_{ij}(x)y^i y^j}$. For a Kropina space $(M, F = \alpha^2/\beta)$, where $\beta = b_i(x)y^i$, we define a new Riemannian metric $h = \sqrt{h_{ij}(x)y^i y^j}$ and a vector field $W = W^i(\partial/\partial x^i)$ of constant length 1 by (1.2) and (1.3). Then, the Kropina metric F satisfies the equation (1.4).*

Conversely, suppose that $h = \sqrt{h_{ij}(x)y^i y^j}$ is a Riemannian metric and $W = W^i(\partial/\partial x^i)$ is a vector field of constant length 1 on (M, h). Consider the metric F defined by (1.4). Then, defining $a_{ij}(x) := e^{-\kappa(x)}h_{ij}(x)$ and $b_i(x) := 2e^{-\kappa(x)}W_i$ by (1.2) using some function $\kappa(x)$ of (x^i) alone, we get $F = \alpha^2/\beta$ and it follows the function $\kappa(x)$ satisfies (1.3).

2 The coefficients of the geodesic spray in a Kropina space.

From the theory of Riemannian spaces, we have the following theorem:

Theorem A([6]) Let (M, g) and $(M, g^* = e^\rho g)$, where $g = \sqrt{g_{ij}(x)y^i y^j}$ and $g^* = \sqrt{g^*_{ij}(x)y^i y^j}$ respectively, be two n-dimensional Riemannian spaces which are comformal each other. Furthermore, let $\gamma_j{}^i{}_k$ and $\gamma^*_j{}^i{}_k$ be the coefficients of Levi-Civita connection of (M, g) and (M, g^*), respectively. Then, we have

$$g^*_{ij} = e^{2\rho}g_{ij}, \quad g^{*ij} = e^{-2\rho}g^{ij}, \quad \gamma^*_j{}^i{}_k = \gamma_j{}^i{}_k + \rho_j\delta^i{}_k + \rho_k\delta^i{}_j - \rho^i g_{jk},$$

where $\rho_i = \partial\rho/\partial x^i$ and $\rho^i = g^{ij}\rho_j$.

From (1.2), we have $h_{ij} = e^\kappa a_{ij}$. Applying Theorem A , we get

$$ {}^h\gamma_j{}^i{}_k = {}^\alpha\gamma_j{}^i{}_k + \frac{1}{2}\kappa_j\delta^i{}_k + \frac{1}{2}\kappa_k\delta^i{}_j - \frac{1}{2}\kappa^i a_{jk}, \tag{2.1}$$

where ${}^h\gamma_j{}^i{}_k$ and ${}^\alpha\gamma_j{}^i{}_k$ are the coefficients of Levi-Civita connection of (M, h) and (M, α) respectively, $\kappa_i = \partial\kappa/\partial x^i$ and $\kappa^i = a^{ij}\kappa_j$. Transvecting (2.1) by $y^j y^k$, we get

$$ {}^h\gamma_0{}^i{}_0 = {}^\alpha\gamma_0{}^i{}_0 + \kappa_0 y^i - \frac{1}{2}h_{00}\overline{\kappa}^i, \tag{2.2}$$

where $\overline{\kappa}^i = h^{ij}\kappa_j$.

We denote the covariant derivative in the Riemannian space (M, α) by $({}_{;i})$ and put as follows:

$$s_{ij} := \frac{b_{i;j} - b_{j;i}}{2}, \quad r_{ij} := \frac{b_{i;j} + b_{j;i}}{2}, \quad s_j := b^i s_{ij}, \quad r_j := b^i r_{ij}.$$

In [1], the authors have shown that the coefficients G^i of the geodesic spray in a Finsler space $(M, F = \alpha\phi(\beta/\alpha))$ are given by

$$2G^i = {}^\alpha\gamma_0{}^i{}_0 + 2\omega\alpha s^i{}_0 + 2\Theta(r_{00} - 2\alpha\omega s_0)\Big(\frac{y^i}{\alpha} + \frac{\omega'}{\omega - s\omega'}b^i\Big), \tag{2.3}$$

where

$$\omega := \frac{\phi'}{\phi - s\phi'}, \quad \Theta := \frac{\omega - s\omega'}{2\{1 + s\omega + (b^2 - s^2)\omega'\}}.$$

For a Kropina space, we have $\phi(s) = 1/s$, hence we obtain

$$\phi' = -\frac{1}{s^2}, \quad \omega = -\frac{1}{2s}, \quad \omega' = \frac{1}{2s^2}$$

and

$$\frac{\omega'}{\omega - s\omega'} = -\frac{1}{2s}, \;\; r_{00} - 2\alpha\omega s_0 = r_{00} + F s_0, \;\; 1 + s\omega + (b^2 - s^2)\omega' = \frac{b^2}{2s^2}, \;\; \Theta = -\frac{s}{b^2}.$$

Substituting the above equalities in (2.3) and using (2.2), we get

$$2G^i \;=\; {}^h\gamma_0{}^i{}_0 - \kappa_0 y^i + \frac{1}{2}h_{00}\overline{\kappa}^i - F s^i{}_0 - \frac{1}{b^2}(r_{00} + F s_0)(\frac{2}{F}y^i - b^i).$$

From Theorem 1, for a Kropina space $(M, \alpha^2/\beta)$, a new Riemannian metric $h = \sqrt{h_{ij}(x)y^i y^j}$ and a vector field $W = W^i(\partial/\partial x^i)$ are defined by (1.2) and (1.3). So, the vector field W satisfies the condition $|W| = 1$ and we have $F = h_{00}/2W_0$.

Therefore, we get

$$2G^i = {}^h\gamma_0{}^i{}_0 + 2\Phi^i, \tag{2.4}$$

where

$$2\Phi^i := -\kappa_0 y^i + \frac{1}{2}h_{00}\overline{\kappa}^i - \frac{h_{00}}{2W_0}s^i{}_0 - \frac{1}{b^2}(r_{00} + \frac{h_{00}s_0}{2W_0})(\frac{4W_0}{h_{00}}y^i - b^i). \tag{2.5}$$

Using (2.1), we have

$$b_{i;j} = 2e^{-\kappa}W_{i||j} + e^{-\kappa}\Big(\kappa_i W_j - \kappa_j W_i - W_r\overline{\kappa}^r h_{ij}\Big),$$

where the notation $({}_{||i})$ stands for the h-covariant derivative in the Riemannian space (M, h).

Remark 1 *We can introduce a Finsler connection* $\Gamma^* = ({}^h\gamma_j{}^i{}_k(x), N_j{}^i := {}^h\gamma_j{}^i{}_k(x)y^k, C_j{}^i{}_k)$ *associated with the linear connection* ${}^h\gamma_j{}^i{}_k(x)$ *of the Riemannian space* (M, h)*. The* h*-covariant derivative are defined as follows ([2]):*

For a vector field $W^i(x)$ *on* M*,*

$$1, \quad \begin{aligned} W^i(x)_{||j} &= \frac{\partial W^i}{\partial x^j} - \frac{\partial W^i}{\partial y^s}N_j{}^s +{}^h \gamma_j{}^i{}_s W^s \\ &= \frac{\partial W^i}{\partial x^j} +{}^h \gamma_j{}^i{}_s W^s. \end{aligned}$$

For a reference vector y^i*,*

$$2, \quad \begin{aligned} y^i{}_{||j} &= \frac{\partial y^i}{\partial x^j} - \frac{\partial y^i}{\partial y^s}N_j{}^s +{}^h \gamma_j{}^i{}_s y^s \\ &= -N_j{}^i + N_j{}^i \\ &= 0. \end{aligned}$$

We put

$$\mathtt{R}_{ij} := \frac{W_{i||j} + W_{j||i}}{2}, \quad \mathtt{S}_{ij} := \frac{W_{i||j} - W_{j||i}}{2}, \qquad \mathtt{R}^i{}_j := h^{ir}\mathtt{R}_{rj}, \quad \mathtt{S}^i{}_j := h^{ir}\mathtt{S}_{rj}$$

$$\mathtt{R}_i := W^r\mathtt{R}_{ri}, \quad \mathtt{S}_i := W^r\mathtt{S}_{ri}, \qquad \mathtt{R}^i := h^{ir}\mathtt{R}_r, \quad \mathtt{S}^i := h^{ir}\mathtt{S}_r.$$

It follows

$$r_{ij} = 2e^{-\kappa}\Big(\mathtt{R}_{ij} - \frac{1}{2}W_r\overline{\kappa}^r h_{ij}\Big), \quad s_{ij} = 2e^{-\kappa}\Big(\mathtt{S}_{ij} + \frac{\kappa_i W_j - \kappa_j W_i}{2}\Big).$$

Furthermore, we get

$$\begin{aligned} s^i{}_j &= 2\mathtt{S}^i{}_j + \overline{\kappa}^i W_j - \kappa_j W^i, & s^i{}_0 &= 2\mathtt{S}^i{}_0 + W_0\overline{\kappa}^i - \kappa_0 W^i \\ s_i &= 2e^{-\kappa}\Big(2\mathtt{S}_i + W_r\overline{\kappa}^r W_i - \kappa_i\Big), & s_0 &= 2e^{-\kappa}\Big(2\mathtt{S}_0 + W_r\overline{\kappa}^r W_0 - \kappa_0\Big), \\ r_{00} &= 2e^{-\kappa}\Big(\mathtt{R}_{00} - \frac{1}{2}W_r\overline{\kappa}^r h_{00}\Big) & b^i &= a^{ir}b_r = e^{\kappa}h^{ir}\frac{2W_r}{e^{\kappa}} = 2W^i. \end{aligned}$$

Substituting the above equality in (2.5), we have

$$2\Phi^i = \frac{h_{00}}{W_0}(\mathtt{S}_0 W^i - \mathtt{S}^i{}_0) + (\mathtt{R}_{00}W^i - 2\mathtt{S}_0 y^i) - \frac{2W_0}{h_{00}}\mathtt{R}_{00}y^i. \tag{2.6}$$

Multiplying now the above equality by $2h_{00}W_0$, we get

$$4h_{00}W_0\Phi^i = 2(h_{00})^2(\mathtt{S}_0W^i - \mathtt{S}^i{}_0) + 2h_{00}W_0(\mathtt{R}_{00}W^i - 2\mathtt{S}_0y^i) - 4(W_0)^2\mathtt{R}_{00}y^i$$

and by putting

$$A^i_{(1)} := 2(\mathtt{S}_0W^i - \mathtt{S}^i{}_0), \quad A^i_{(2)} := 2(\mathtt{R}_{00}W^i - 2\mathtt{S}_0y^i), \quad A^i_{(3)} := -4\mathtt{R}_{00}y^i,$$

it follows

$$4h_{00}W_0\Phi^i = (h_{00})^2A^i_{(1)} + h_{00}W_0A^i_{(2)} + (W_0)^2A^i_{(3)}. \tag{2.7}$$

3 The necessary and sufficient conditions for a Kropina space to be of constant curvature.

In this section, we consider a Kropina space $(M, F = \alpha^2/\beta)$ of constant curvatue K, where $\alpha = \sqrt{a_{ij}(x)y^iy^j}$ and $\beta = b_i(x)y^i$. Furthermore, we suppose that the matrix (a_{ij}) is always positive definite and that the dimension n is more than or equal two. Hence, it follows that α^2 is not divisible by β. This is an important relation and is equivalent to that h_{00} is not divisible by W_0. Using these, we will obtain the necessary and sufficient conditions for a Kropina space to be of constant curvature.

3.1 The curvature tensor of a Kropina space.

Let $R_j{}^i{}_{kl}$ be the h-curvature tensor of Cartan connection in Finsler space. The Berwald's spray curvature tensor is

$${}^{(b)}R_j{}^i{}_{kl} \;=\; A_{(kl)}\Big\{\frac{\partial G_j{}^i{}_k}{\partial x^l} + G_j{}^r{}_kG_r{}^i{}_l\Big\}. \tag{3.1}$$

It is well-known that the equality $R_0{}^i{}_{kl} ={}^{(b)} R_0{}^i{}_{kl}$ holds good ([2]).

From $2G^i = {}^h\gamma_0{}^i{}_0 + 2\Phi^i$, it follows

$$G^i{}_j = {}^h\gamma_0{}^i{}_j + \Phi^i{}_j \quad and \quad G_j{}^i{}_k = {}^h\gamma_j{}^i{}_k + \Phi_j{}^i{}_k.$$

Substituting the above equalities in (3.1), we get

$$R_j{}^i{}_{kl} = {}^hR_j{}^i{}_{kl} + A_{(kl)}\Big\{\Phi_j{}^i{}_{k||l} + \Phi_j{}^r{}_k\Phi_r{}^i{}_l\Big\}.$$

The following Proposition is well-known ([2]):

Proposition 1 *The necessary and sufficient condition for a Finsler space (M, F) to be of scalar curvature K is that the equality*

$$R_0{}^i{}_{0l} = KF^2(\delta^i{}_l - l^i l_l), \tag{3.2}$$

where $l^i = y^i/F$ and $l_l = \partial F/\partial y^l$, holds.

If the equality (3.2) holds good and K is constant, then the Finsler space is called of *constant curvature K*.

For a Kropina space of constant curvatue K, the equality (3.2) holds and K is constant.

Since $F = \alpha^2/\beta$ is written as $F = h_{00}/(2W_0)$, we have

$$\delta^i{}_l - l^i l_l = \delta^i{}_l - \frac{2W_0h_{0l} - h_{00}W_l}{h_{00}W_0}y^i.$$

Using the curvature we obtained above, we have

$$R_0{}^i{}_{0l} = {}^hR_0{}^i{}_{0l} + 2\Phi^i{}_{||l} - \Phi^i{}_{l||0} + 2\Phi^r\Phi_r{}^i{}_l - \Phi^r{}_l\Phi^i{}_r.$$

Substituting the above equalities in (3.2), we get

$$(3.3) KF^2\Big(\delta^i{}_l - \frac{2W_0h_{0l} - h_{00}W_l}{h_{00}W_0}y^i\Big) = {}^hR_0{}^i{}_{0l} + 2\Phi^i{}_{||l} - \Phi^i{}_{l||0} + 2\Phi^r\Phi_r{}^i{}_l - \Phi^r{}_l\Phi^i{}_r.$$

3.2 Rewriting the equation (3.3) using h_{00} and W_0.

(1)The calculations for $\Phi^i{}_{||l}$.

First, applying the h-covariant derivative ${}_{||l}$ to (2.7), it follows

$$\begin{aligned} & 4h_{00}W_{0||l}\Phi^i + 4h_{00}W_0\Phi^i{}_{||l} \\ = \ & (h_{00})^2A^i_{(1)_{||l}} + h_{00}W_0A^i_{(2)_{||l}} + h_{00}W_{0||l}A^i_{(2)} + (W_0)^2A^i_{(3)_{||l}} + 2W_0W_{0||l}A^i_{(3)} \end{aligned}$$

and using again (2.7), we get

$$\begin{aligned} & 4h_{00}(W_0)^2\Phi^i{}_{||l} \\ = \ & (h_{00})^2W_0A^i_{(1)_{||l}} - (h_{00})^2A^i_{(1)}W_{0||l} + h_{00}(W_0)^2A^i_{(2)_{||l}} \\ & \qquad + (W_0)^3A^i_{(3)_{||l}} + (W_0)^2A^i_{(3)}W_{0||l}. \end{aligned}$$

Putting now

$$B^i_{(1)l} = A^i_{(1)_{||l}}, \ B^i_{(21)l} = -A^i_{(1)}W_{0||l}, \ B^i_{(22)l} = A^i_{(2)_{||l}}, \ B^i_{(3)l} = A^i_{(3)_{||l}}, \ B^i_{(4)l} = A^i_{(3)}W_{0||l},$$

we have

$$(3.4)\qquad 4h_{00}(W_0)^2\Phi^i{}_{||l} = (h_{00})^2 W_0 B^i_{(1)l} + (h_{00})^2 B^i_{(21)l} + h_{00}(W_0)^2 B^i_{(22)l} + (W_0)^3 B^i_{(3)l} + (W_0)^2 B^i_{(4)l}.$$

(2)The calculations for $\Phi^i{}_l$.

Secondly, derivating (2.7) by y^l, we get

$$\begin{aligned}
& 8h_{0l}W_0\Phi^i + 4h_{00}W_l\Phi^i + 4h_{00}W_0\Phi^i{}_l \\
= \ & (h_{00})^2 A^i_{(1)}{}_{\cdot l} + h_{00}W_0 A^i_{(2).l} + h_{00}(4h_{0l}A^i_{(1)} + W_l A^i_{(2)}) \\
& + (W_0)^2 A^i_{(3)}{}_{\cdot l} + 2W_0(W_l A^i_{(3)} + h_{0l}A^i_{(2)}),
\end{aligned}$$

where the notation $({}_{.l})$ stands for the derivation by y^l.

Using the above equality and (2.7), we obtain

$$\begin{aligned}
(3.5)\qquad & 4(h_{00})^2(W_0)^2\Phi^i{}_l \\
= \ & (h_{00})^3 W_0 C^i_{(0)l} + (h_{00})^3 C^i_{(11)l} + (h_{00})^2(W_0)^2 C^i_{(12)l} \\
& + (h_{00})^2 W_0 C^i_{(21)l} + h_{00}(W_0)^3 C^i_{(22)l} + h_{00}(W_0)^2 C^i_{(3)l} + (W_0)^3 C^i_{(4)l},
\end{aligned}$$

where

$$\begin{aligned}
& C^i_{(0)l} := A^i_{(1)}{}_{\cdot l} = 2(\mathtt{S}_l W^i - \mathtt{S}^i{}_l), \ \ C^i_{(11)l} := -A^i_{(1)}W_l = -2(\mathtt{S}_0 W^i - \mathtt{S}^i{}_0)W_l, \\
& C^i_{(12)l} := A^i_{(2).l} = 4(\mathtt{R}_{0l}W^i - \mathtt{S}_l y^i - \mathtt{S}_0\delta^i{}_l), \ \ C^i_{(21)l} := 2A^i_{(1)}h_{0l} = 4(\mathtt{S}_0 W^i - \mathtt{S}^i{}_0)h_{0l}, \\
& C^i_{(22)l} := A^i_{(3)}{}_{\cdot l} = -4(2\mathtt{R}_{0l}y^i + \mathtt{R}_{00}\delta^i{}_l), \ \ C^i_{(3)l} := W_l A^i_{(3)} = -4\mathtt{R}_{00}W_l y^i, \\
& C^i_{(4)l} := -2A^i_{(3)}h_{0l} = 8\mathtt{R}_{00}h_{0l}y^i.
\end{aligned}$$

(3)The calculations for $\Phi^i{}_{l||0}$.

Applying the h-covariant derivaive ${}_{||0}$ to (3.5), we get

$$\begin{aligned}
& 8(h_{00})^2 W_0 W_{0||0}\Phi^i{}_l + 4(h_{00})^2(W_0)^2\Phi^i{}_{l||0} \\
= \ & (h_{00})^3 W_0 C^i_{(0)l||0} + (h_{00})^3(W_{0||0}C^i_{(0)l} + C^i_{(11)l||0}) \\
& + (h_{00})^2(W_0)^2 C^i_{(12)l||0} + (h_{00})^2 W_0(2W_{0||0}C^i_{(12)l} + C^i_{(21)l||0}) \\
& + (h_{00})^2 W_{0||0}C^i_{(21)l} + h_{00}(W_0)^3 C^i_{(22)l||0} + h_{00}(W_0)^2(3W_{0||0}C^i_{(22)l} + C^i_{(3)l||0}) \\
& + 2h_{00}W_0W_{0||0}C^i_{(3)l} + (W_0)^3 C^i_{(4)l||0} + 3(W_0)^2 W_{0||0}C^i_{(4)l}.
\end{aligned}$$

Using the above equality and (3.5), we obtain

$$\begin{aligned}
(3.6)\qquad & 4(h_{00})^2(W_0)^3\Phi^i{}_{l||0} \\
= \ & (h_{00})^3(W_0)^2 D^i_{(1)l} + (h_{00})^3 W_0 D^i_{(21)l} + (h_{00})^3 D^i_{(31)l} \\
& + (h_{00})^2(W_0)^3 D^i_{(22)l} + (h_{00})^2(W_0)^2 D^i_{(32)l} + (h_{00})^2 W_0 D^i_{(41)l} \\
& + h_{00}(W_0)^4 D^i_{(33)l} + h_{00}(W_0)^3 D^i_{(42)l} + (W_0)^4 D^i_{(5)l} + (W_0)^3 D^i_{(6)l},
\end{aligned}$$

where

$$\begin{aligned}
& D^i_{(1)l} = C^i_{(0)l||0}, \quad D^i_{(21)l} = C^i_{(11)l||0} - W_{0||0}C^i_{(0)l}, \quad D^i_{(31)l} = -2W_{0||0}C^i_{(11)l}, \\
& D^i_{(22)l} = C^i_{(12)l||0}, \quad D^i_{(32)l} = C^i_{(21)l||0}, \quad D^i_{(41)l} = -W_{0|0}C^i_{(21)l}, \\
& D^i_{(33)l} = C^i_{(22)l||0}, \quad D^i_{(42)l} = W_{0||0}C^i_{(22)l} + C^i_{(3)l||0}, \quad D^i_{(5)l} = C^i_{(4)l||0}, \quad D^i_{(6)l} = W_{0||0}C^i_{(4)l}.
\end{aligned}$$

(4)The calculations for $\Phi^r{}_l\Phi^i{}_r$.

Using (3.5), we have

$$
\begin{aligned}
(3.7)\qquad & 16(h_{00})^4(W_0)^4\Phi^r{}_l\Phi^i{}_r \\
= \; & (h_{00})^6(W_0)^2E^i_{(01)l} + (h_{00})^6W_0E^i_{(11)l} + (h_{00})^6E^i_{(21)l} + (h_{00})^5(W_0)^3E^i_{(12)l} \\
& +(h_{00})^5(W_0)^2E^i_{(22)l} + (h_{00})^5W_0E^i_{(31)l} + (h_{00})^4(W_0)^4E^i_{(23)l} + (h_{00})^4(W_0)^3E^i_{(32)l} \\
& +(h_{00})^4(W_0)^2E^i_{(41)l} + (h_{00})^3(W_0)^5E^i_{(33)l} + (h_{00})^3(W_0)^4E^i_{(42)l} + (h_{00})^3(W_0)^3E^i_{(51)l} \\
& +(h_{00})^2(W_0)^6E^i_{(43)l} + (h_{00})^2(W_0)^5E^i_{(52)l} + (h_{00})^2(W_0)^4E^i_{(61)l} + h_{00}(W_0)^6E^i_{(62)l} \\
& +h_{00}(W_0)^5E^i_{(7)l} + (W_0)^6E^i_{(8)l},
\end{aligned}
$$

where

$$
\begin{aligned}
E^i_{(0)l} : &= C^i_{(0)r}C^r_{(0)l}, \quad E^i_{(11)l} := C^i_{(11)r}C^r_{(0)l} + C^i_{(0)r}C^r_{(11)l}, \quad E^i_{(21)l} := C^i_{(11)r}C^r_{(11)l}, \\
E^i_{(12)l} : &= C^i_{(0)r}C^r_{(12)l} + C^i_{(12)r}C^r_{(0)l}, \\
E^i_{(22)l} : &= C^i_{(12)r}C^r_{(11)l} + C^i_{(11)r}C^r_{(12)l} + C^i_{(21)r}C^r_{(0)l} + C^i_{(0)r}C^r_{(21)l}, \\
E^i_{(31)l} : &= C^i_{(21)r}C^r_{(11)l} + C^i_{(11)r}C^r_{(21)l}, \quad E^i_{(23)l} := C^i_{(12)r}C^r_{(12)l} + C^i_{(22)r}C^r_{(0)l} + C^i_{(0)r}C^r_{(22)l}, \\
E^i_{(32)l} : &= C^i_{(21)r}C^r_{(12)l} + C^i_{(12)r}C^r_{(21)l} + C^i_{(3)r}C^r_{(0)l} + C^i_{(22)r}C^r_{(11)l} + C^i_{(11)r}C^r_{(22)l} + C^i_{(0)r}C^r_{(3)l} \\
E^i_{(41)l} : &= C^i_{(3)r}C^r_{(11)l} + C^i_{(11)r}C^r_{(3)l} + C^i_{(21)r}C^r_{(21)l}, \quad E^i_{(33)l} := C^i_{(22)r}C^r_{(12)l} + C^i_{(12)r}C^r_{(22)l}, \\
E^i_{(42)l} : &= C^i_{(4)r}C^r_{(0)l} + C^i_{(3)r}C^r_{(12)l} + C^i_{(22)r}C^r_{(21)l} + C^i_{(21)r}C^r_{(22)l} + C^i_{(12)r}C^r_{(3)l} + C^i_{(0)r}C^r_{(4)l}, \\
E^i_{(51)l} : &= C^i_{(3)r}C^r_{(21)l} + C^i_{(21)r}C^r_{(3)l} + C^i_{(4)r}C^r_{(11)l} + C^i_{(11)r}C^r_{(4)l}, \quad E^i_{(43)l} := C^i_{(22)r}C^r_{(22)l}, \\
E^i_{(52)l} : &= C^i_{(4)r}C^r_{(12)l} + C^i_{(12)r}C^r_{(4)l} + C^i_{(3)r}C^r_{(22)l} + C^i_{(22)r}C^r_{(3)l}, \\
E^i_{(61)l} : &= C^i_{(3)r}C^r_{(3)l} + C^i_{(4)r}C^r_{(21)l} + C^i_{(21)r}C^r_{(4)l}, \quad E^i_{(62)l} := C^i_{(4)r}C^r_{(22)l} + C^i_{(22)r}C^r_{(4)l}, \\
E^i_{(7)l} : &= C^i_{(4)r}C^r_{(3)l} + C^i_{(3)r}C^r_{(4)l}, \quad E^i_{(8)l} := C^i_{(4)r}C^r_{(4)l}.
\end{aligned}
$$

(5)The calculations for $\Phi^r\Phi_r{}^i{}_l$.
Derivating (3.5) by y^r, we get

$$
\begin{aligned}
& 16h_{00}h_{0r}(W_0)^2\Phi^i{}_l + 8(h_{00})^2W_0W_r\Phi^i{}_l + 4(h_{00})^2(W_0)^2\Phi_r{}^i{}_l \\
= \; & 6(h_{00})^2h_{0r}W_0C^i_{(0)l} + (h_{00})^3W_rC^i_{(0)l} + (h_{00})^3W_0C^i_{(0)l\cdot r} \\
& +6(h_{00})^2h_{0r}C^i_{(11)l} + (h_{00})^3C^i_{(11)l\cdot r} \\
& +4h_{00}h_{0r}(W_0)^2C^i_{(12)l} + 2(h_{00})^2W_0W_rC^i_{(12)l} + (h_{00})^2(W_0)^2C^i_{(12)l\cdot r} \\
& +4h_{00}h_{0r}W_0C^i_{(21)l} + (h_{00})^2W_rC^i_{(21)l} + (h_{00})^2W_0C^i_{(21)l\cdot r} \\
& +2h_{0r}(W_0)^3C^i_{(22)l} + 3h_{00}(W_0)^2W_rC^i_{(22)l} + h_{00}(W_0)^3C^i_{(22)l\cdot r} \\
& +2h_{0r}(W_0)^2C^i_{(3)l} + 2h_{00}W_0W_rC^i_{(3)l} + h_{00}(W_0)^2C^i_{(3)l\cdot r} \\
& +3(W_0)^2W_rC^i_{(4)l} + (W_0)^3C^i_{(4)l\cdot r}.
\end{aligned}
$$

Using the above equality, (3.5) and $C^i_{(0)l\cdot r} = 0$, we have

$$
\begin{aligned}
& 4(h_{00})^3(W_0)^3\Phi_r{}^i{}_l \\
= \; & (h_{00})^4W_0H^i_{(01)rl} + (h_{00})^4H^i_{(11)rl} + (h_{00})^3(W_0)^3H^i_{(02)rl} + (h_{00})^3(W_0)^2H^i_{(12)rl} \\
& +(h_{00})^3W_0H^i_{(21)rl} + (h_{00})^2(W_0)^4H^i_{(13)rl} + (h_{00})^2(W_0)^3H^i_{(22)rl} \\
& +h_{00}(W_0)^4H^i_{(3)rl} + h_{00}(W_0)^3H^i_{(4)rl} + (W_0)^4H^i_{(5)rl},
\end{aligned}
$$

where

$$
H^i_{(01)rl} = C^i_{(11)l\cdot r} - W_rC^i_{(0)l}, \quad H^i_{(11)rl} = -2W_rC^i_{(11)l}, \quad H^i_{(02)rl} = C^i_{(12)l.r},
$$

$$\begin{aligned}
&H^i_{(12)rl} = 2h_{0r}C^i_{(0)l} + C^i_{(21)l\cdot r}, \quad H^i_{(21)rl} = -W_rC^i_{(21)l} + 2h_{0r}C^i_{(11)l},\\
&H^i_{(13)rl} = C^i_{(22)l\cdot r}, \quad H^i_{(22)rl} = W_rC^i_{(22)l} + C^i_{(3)l\cdot r}, \quad H^i_{(3)rl} = C^i_{(4)l\cdot r} - 2h_{0r}C^i_{(22)l},\\
&H^i_{(4)rl} = W_rC^i_{(4)l} - 2h_{0r}C^i_{(3)l}, \quad H^i_{(5)rl} = -4h_{0r}C^i_{(4)l}.
\end{aligned}$$

Using the above equality and (2.7), we get

$$\begin{aligned}
&(3.8) 16(h_{00})^4(W_0)^4\Phi^r\Phi_r{}^i{}_l\\
= \ &(h_{00})^6W_0J^i_{(11)l} + (h_{00})^6J^i_{(21)l} + (h_{00})^5(W_0)^3J^i_{(12)l} + (h_{00})^5(W_0)^2J^i_{(22)l}\\
&+(h_{00})^5W_0J^i_{(31)l} + (h_{00})^4(W_0)^4J^i_{(23)l} + (h_{00})^4(W_0)^3J^i_{(32)l} + (h_{00})^4(W_0)^2J^i_{(41)l}\\
&+(h_{00})^3(W_0)^5J^i_{(33)l} + (h_{00})^3(W_0)^4J^i_{(42)l} + (h_{00})^3(W_0)^3J^i_{(51)l} + (h_{00})^2(W_0)^6J^i_{(43)l}\\
&+(h_{00})^2(W_0)^5J^i_{(52)l} + (h_{00})^2(W_0)^4J^i_{(61)l} + h_{00}(W_0)^6J^i_{(62)l} + h_{00}(W_0)^5J^i_{(71)l} + (W_0)^6J^i_{(8)l},
\end{aligned}$$

where

$$\begin{aligned}
&J^i_{(11)l} = A^r_{(1)}H^i_{(01)rl}, \quad J^i_{(21)l} = A^r_{(1)}H^i_{(11)rl}, \quad J^i_{(12)l} = A^r_{(1)}H^i_{(02)rl},\\
&J^i_{(22)l} = A^r_{(1)}H^i_{(12)rl} + A^r_{(2)}H^i_{(01)rl}, \quad J^i_{(31)l} = A^r_{(1)}H^i_{(21)rl} + A^r_{(2)}H^i_{(11)rl},\\
&J^i_{(23)l} = A^r_{(1)}H^i_{(13)rl} + A^r_{(2)}H^i_{(02)rl}, \quad J^i_{(32)l} = A^r_{(1)}H^i_{(22)rl} + A^r_{(2)}H^i_{(12)rl} + A^r_{(3)}H^i_{(01)rl},\\
&J^i_{(41)l} = A^r_{(2)}H^i_{(21)rl} + A^r_{(3)}H^i_{(11)rl}, \quad J^i_{(33)l} = A^r_{(2)}H^i_{(13)rl} + A^r_{(3)}H^i_{(02)rl},\\
&J^i_{(42)l} = A^r_{(1)}H^i_{(3)rl} + A^r_{(2)}H^i_{(22)rl} + A^r_{(3)}H^i_{(12)rl},\\
&J^i_{(51)l} = A^r_{(1)}H^i_{(4)rl} + A^r_{(3)}H^i_{(21)rl}, \quad J^i_{(43)l} = A^r_{(3)}H^i_{(13)rl},\\
&J^i_{(52)l} = A^r_{(2)}H^i_{(3)rl} + A^r_{(3)}H^i_{(22)rl}, \quad J^i_{(61)l} = A^r_{(2)}H^i_{(4)rl} + A^r_{(1)}H^i_{(5)rl},\\
&J^i_{(62)l} = A^r_{(3)}H^i_{(3)rl}, \quad J^i_{(7)l} = A^r_{(2)}H^i_{(5)rl} + A^r_{(3)}H^i_{(4)rl}, \quad J^i_{(8)l} = A^r_{(3)}H^i_{(5)rl}.
\end{aligned}$$

(6)The main relation.

Multiplying (3.3) by $16(h_{00})^4(W_0)^4$ and using $F^2 = (h_{00})^2/\{4(W_0)^2\}$, we have the equality

$$\begin{aligned}
&4K(h_{00})^6(W_0)^2h^i{}_l = 16(h_{00})^4(W_0)^4\cdot {}^hR_0{}^i{}_{0l} + 8(h_{00})^3(W_0)^2\cdot 4h_{00}(W_0)^2\Phi^i{}_{||l}\\
&-4(h_{00})^2W_0\cdot 4(h_{00})^2(W_0)^3\Phi^i{}_{l||0} + 2\cdot 16(h_{00})^4(W_0)^4\Phi^r\Phi_r{}^i{}_l - 16(h_{00})^4(W_0)^4\Phi^r{}_l\Phi^i{}_r.
\end{aligned}$$

Substituting (3.4), (3.5), (3.6), (3.7) and (3.8) in the above equality, we get

$$\begin{aligned}
&4K(h_{00})^6(W_0)^2\delta^i{}_l - 8K(h_{00})^5(W_0)^2h_{0l}y^i + 4K(h_{00})^6W_0W_ly^i\\
= \ &-(h_{00})^6(W_0)^2E^i_{(0)l} + (h_{00})^6W_0(2J^i_{(11)l} - E^i_{(11)l})\\
&+(h_{00})^6(2J^i_{(21)l} - E^i_{(21)l}) + (h_{00})^5(W_0)^3(8B^i_{(1)l} - 4D^i_{(1)l} + 2J^i_{(12)l} - E^i_{(12)l})\\
&+(h_{00})^5(W_0)^2(-4D^i_{(21)l} + 8B^i_{(21)l} + 2J^i_{(22)l} - E^i_{(22)l})\\
&+(h_{00})^5W_0(2J^i_{(31)l} - 4D^i_{(31)l} - E^i_{(31)l})\\
&+(h_{00})^4(W_0)^4(16{}^hR_0{}^i{}_{0l} + 8B^i_{(22)l} - 4D^i_{(22)l} + 2J^i_{(23)l} - E^i_{(23)l})\\
&+(h_{00})^4(W_0)^3(2J^i_{(32)l} - 4D^i_{(32)l} - E^i_{(32)l}) + (h_{00})^4(W_0)^2(2J^i_{(41)l} - E^i_{(41)l} - 4D^i_{(41)l})\\
&+(h_{00})^3(W_0)^5(8B^i_{(3)l} - 4D^i_{(33)l} + 2J^i_{(33)l} - E^i_{(33)l})\\
&+(h_{00})^3(W_0)^4(2J^i_{(42)l} + 8B^i_{(4)l} - 4D^i_{(42)l} - E^i_{(42)l})\\
&+(h_{00})^3(W_0)^3(2J^i_{(51)l} - E^i_{(51)l}) + (h_{00})^2(W_0)^6(2J^i_{(43)l} - E^i_{(43)l})\\
&+(h_{00})^2(W_0)^5(2J^i_{(52)l} - 4D^i_{(5)l} - E^i_{(52)l}) + (h_{00})^2(W_0)^4(2J^i_{(61)l} - 4D^i_{(6)l} - E^i_{(61)l})\\
&+h_{00}(W_0)^6(2J^i_{(62)l} - E^i_{(62)l}) + h_{00}(W_0)^5(2J^i_{(71)l} - E^i_{(7)l}) + (W_0)^6(2J^i_{(8)l} - E^i_{(8)l}).
\end{aligned}$$

Taking into account the equalities $J^i_{(21)l} = 0$ and $E^i_{(21)l} = 0$, we have

$$(h_{00})^4 P^i_{(5)_l} + (h_{00})^2 Q^i_{(9)_l} + (W_0)^4 R^i_{(9)_l} = 0, \tag{3.9}$$

where

$$\begin{aligned}
P^i_{(5)_l} &= (h_{00})^2 W_0 (-E^i_{(0)l} - 4K\delta^i{}_l) \\
&\quad +(h_{00})^2 (2J^i_{(11)l} - E^i_{(11)l} - 4KW_l y^i) \\
&\quad +h_{00}(W_0)^2 (8B^i_{(1)l} - 4D^i_{(1)l} + 2J^i_{(12)l} - E^i_{(12)l}) \\
&\quad +h_{00}W_0(-4D^i_{(21)l} + 8B^i_{(21)l} + 2J^i_{(22)l} - E^i_{(22)l} + 8Kh_{0l}y^i) \\
&\quad +(W_0)^3(16{}^hR_0{}^i{}_{0l} + 8B^i_{(22)l} - 4D^i_{(22)l} + 2J^i_{(23)l} - E^i_{(23)l}) \\
&\quad +(W_0)^2(2J^i_{(32)l} - 4D^i_{(32)l} - E^i_{(32)l}), \\
Q^i_{(9)_l} &= (h_{00})^3(2J^i_{(31)l} - 4D^i_{(31)l} - E^i_{(31)l}) \\
&\quad +(h_{00})^2 W_0(2J^i_{(41)l} - E^i_{(41)l} - 4D^i_{(41)l}) \\
&\quad +h_{00}(W_0)^4(8B^i_{(3)l} - 4D^i_{(33)l} + 2J^i_{(33)l} - E^i_{(33)l}) \\
&\quad +h_{00}(W_0)^3(2J^i_{(42)l} + 8B^i_{(4)l} - 4D^i_{(42)l} - E^i_{(42)l}) \\
&\quad +h_{00}(W_0)^2(2J^i_{(51)l} - E^i_{(51)l}) \\
&\quad +(W_0)^5(2J^i_{(43)l} - E^i_{(43)l}) \\
&\quad +(W_0)^4(2J^i_{(52)l} - 4D^i_{(5)l} - E^i_{(52)l}) \\
&\quad +(W_0)^3(2J^i_{(61)l} - 4D^i_{(6)l} - E^i_{(61)l}) \\
R^i_{(9)_l} &= h_{00}W_0(2J^i_{(62)l} - E^i_{(62)l}) \\
&\quad +h_{00}(2J^i_{(7)l} - E^i_{(7)l}) \\
&\quad +W_0(2J^i_{(8)l} - E^i_{(8)l}).
\end{aligned}$$

We call $P^i_{(5)_l}$, $Q^i_{(9)_l}$ and $R^i_{(9)_l}$ the *curvature part*, the *vanishing part* and the *Killing part*, respectively.

Proposition 2 *The necessary and sufficient condition for a Kropina space* $(M, F = \alpha^2/\beta = h_{00}/2W_0)$ *to be of constant curvature* K *is that (3.9) holds good.*

3.3 The Killing part.

We consider the Killing part $R^i_{(9)_l}$ and obtain the conclusion that the vector field W is Killing.

First, we have

$$\begin{aligned}
J^i_{(62)l} &= -64\mathtt{R}_{00}(2\mathtt{R}_{00}h_{0l} + h_{00}\mathtt{R}_{0l})y^i - 32(\mathtt{R}_{00})^2 h_{00}\delta^i{}_l, \\
E^i_{(62)l} &= -64\mathtt{R}_{00}h_{00}\mathtt{R}_{0l}y^i - 128(\mathtt{R}_{00})^2 h_{0l}y^i, \\
J^i_{(7)l} &= -32\mathtt{R}_{00}\{(3\mathtt{R}_{00}W_0 - 4\mathtt{S}_0 h_{00})h_{0l} + h_{00}\mathtt{R}_{00}W_l\}y^i, \\
E^i_{(7)l} &= -32(\mathtt{R}_{00})^2(h_{00}W_l y^i + W_0 h_{0l}y^i), \\
2J^i_{(8)l} - E^i_{(8)l} &= 192(\mathtt{R}_{00})^2 h_{00}h_{0l}y^i.
\end{aligned}$$

Using the above equalities, we get

$$R^i_{(9)_l} = -32h_{00}\mathtt{R}_{00}\Big(W_0(2\mathtt{R}_{00}h_{00}\delta^i{}_l + 2h_{00}\mathtt{R}_{0l}y^i + 7\mathtt{R}_{00}h_{0l}y^i) - 8\mathtt{S}_0 h_{00}h_{0l}y^i + \mathtt{R}_{00}h_{00}W_l y^i\Big).$$

Substituting the above equality in (3.9) and dividing it by $W_0 h_{00}$, we get

$$\begin{aligned}(3.10)\qquad &(h_{00})^3 P^i_{(5)l} + h_{00} Q^i_{(9)l}\\ &\quad -32(W_0)^4 \mathtt{R}_{00}\Big(W_0(2\mathtt{R}_{00} h_{00} \delta^i{}_l + 2h_{00} \mathtt{R}_{0l} y^i + 7\mathtt{R}_{00} h_{0l} y^i)\\ &\qquad\qquad -8\mathtt{S}_0 h_{00} h_{0l} y^i + \mathtt{R}_{00} h_{00} W_l y^i \Big) = 0.\end{aligned}$$

Lemma 1 *In the equation (3.10), it follows that* $\mathtt{R}_{00}$ *is divisible by* h_{00}.

(*Proof.*) Suppose that $\mathtt{R}_{00}$ is not divisible by h_{00} and since (h_{ij}) is positive definite, $(R_{00})^2$ is not divisible by h_{00}.

Taking into account that $P^i_{(5)l}$ and $Q^i_{(9)l}$ are homogeneous polynomials of y^i and that $(W_0)^2$ is not divisible by h_{00}, it follows that the equation

$$h_{0l} y^i = h_{00} \eta^i{}_l,$$

where $\eta_l{}^i(x)$ is a function of (x^i) alone, holds good. Transvecting the above equation by W^l, we get

$$W_0 y^i = h_{00} \eta_l{}^i(x) W^l.$$

Since h_{00} is not divisible by W_0, the above equation is impossible. Q.E.D.

Therefore, it follows that $\mathtt{R}_{00}$ is divisible by h_{00} and the following equation holds:

$$\mathtt{R}_{00} = c(x) h_{00},$$

where $c(x)$ is a function of (x^i) alone. Derivating the above equation by y^i and y^j, we get

$$(3.11)\qquad W_{i||j} + W_{j||i} = 2c(x) h_{ij}.$$

Transvecting (3.11) by $W^i W^j$, we get $W_{i||j} W^i W^j = c(x) h_{ij} W^i W^j$ and using $h_{ij} W^i W^j = |W|^2 = 1$ and $W_{i||r} W^i = 0$, we obtain $c(x) = 0$. Therefore, it follows that the equality

$$(3.12)\qquad \mathtt{R}_{ij} = 0$$

holds good. Hence, we have that W is a Killing vector field. Therefore, we can state

Lemma 2 *If a Kropina space* $(M, \alpha^2/\beta)$ *is of constant curvature* K*, then*
(1) $W(x)$ *is a Killing vector field,*
and then
(2) Killing part $R^i_{(9)l} = 0$.

Using (3.12), the equation (3.10) reduces to

$$(3.13)\qquad (h_{00})^2 P^i_{(5)l} + Q^i_{(9)l} = 0$$

and we have the following equalities:

$$(3.14)\quad W_{i||j} = \mathtt{S}_{ij},\ \ \mathtt{S}_j = W_{i||j} W^i = 0,\ \ W_{0||j} = \mathtt{S}_{0j},\ \ W_{i||0} = \mathtt{S}_{i0},\ \ W_{0||0} = 0.$$

3.4 The vanishing part.

In this subsection, we will show that the equality $Q^i_{(9)l} = 0$ holds from the conclusion $\mathtt{R}_{00} = 0$ in the previous subsection.

Using (3.12) and (3.14), the $A's$, $B's$, $C's$, $D's$, $E's$, $H's$ and $J's$ reduce to

$$
\begin{aligned}
&A^i_{(1)} := -2W^i{}_{||0},\ \ B^i_{(1)l} = -2W^i{}_{||0||l},\ \ B^i_{(21)l} = 2W^i{}_{||0}W_{0||l},\ \ C^i_{(0)l} = -2W^i{}_{||l},\\
&C^i_{(11)l} = 2W^i{}_{||0}W_l,\ \ C^i_{(21)l} = -4W^i{}_{||0}h_{0l},\ \ D^i_{(1)l} = -2W^i{}_{||l||0},\\
&D^i_{(21)l} = 2W^i{}_{||0||0}W_l + 2W^i{}_{||0}W_{l||0},\ \ D^i_{(32)l} = -4W^i{}_{||0||0}h_{0l},\ \ E^i_{(0)l} = 4W^i{}_{||r}W^r{}_{||l},\\
&E^i_{(11)l} = -4W^i{}_{||r}W^r{}_{||0}W_l,\ \ E^i_{(22)l} = 8W^i{}_{||0}W_{0||l} + 8W^i{}_{||r}W^r{}_{||0}h_{0l},\\
&H^i_{(01)rl} = 2W^i{}_{||r}W_l + 2W_rW^i{}_{||l},\ \ H^i_{(11)rl} = -4W_rW^i{}_{||0}W_l,\\
&H^i_{(12)rl} = -4(h_{0r}W^i{}_{||l} + W^i{}_{||r}h_{0l} + W^i{}_{||0}h_{rl}),\ \ H^i_{(21)rl} = 4(W_rW^i{}_{||0}h_{0l} + h_{0r}W^i{}_{||0}W_l),\\
&J^i_{(11)l} = -4W^i{}_{||r}W^r{}_{||0}W_l,\ \ J^i_{(22)l} = 8W^i{}_{||r}W^r{}_{||0}h_{0l} + 8W^i{}_{||0}W_{l||0},
\end{aligned}
$$

and the others are zero. Using the above equalities, we get

$$
\begin{aligned}
&2J^i_{(31)l} - 4D^i_{(31)l} - E^i_{(31)l} = 0,\quad 2J^i_{(41)l} - E^i_{(41)l} - 4D^i{}_{(41)l} = 0\\
&8B^i_{(3)} - 4D^i_{(33)l} + 2J^i_{(33)l} - E^i_{(33)l} = 0,\quad 2J^i_{(42)l} + 8B^i_{(4)} - 4D^i_{(42)l} - E^i_{(42)l} = 0,\\
&2J^i_{(51)l} - E^i_{(51)l} = 0,\quad 2J^i_{(43)l} - E^i_{(43)l} = 0,\\
&2J^i_{(52)l} - 4D^i_{(5)l} - E^i_{(52)l} = 0,\quad 2J^i_{(61)l} - 4D^i_{(6)l} - E^i_{(61)l} = 0.
\end{aligned}
$$

Therefore, from Lemma 2, it follows

Lemma 3 *If a Kropina space $(M, \alpha^2/\beta)$ is of constant curvature K, then*
(1) vanishing part $Q^i_{(9)l} = 0$,
and then
(2) curvature part $P^i_{(5)l} = 0$.

3.5 The curvature part.

In this subsection, we will show that Lemma 3 implies that (M, h) is a Riemannian space of constant curvature K.

Using the resuls given at the beginning of the previous subsection, we have

$$
\begin{aligned}
-E^i_{(0)l} - 4K\delta^i{}_l &= -4W^i{}_{||r}W^r{}_{||l} - 4K\delta^i{}_l,\\
2J^i_{(11)l} - E^i_{(11)l} - 4KW_ly^i &= -4W^i{}_{||r}W^r{}_{||0}W_l - 4KW_ly^i,\\
8B^i_{(1)l} - 4D^i_{(1)l} + 2J^i_{(12)l} - E^i_{(12)l} &= -16W^i{}_{||0||l} + 8W^i{}_{||l||0},\\
-4D^i_{(21)l} + 8B^i_{(21)l} + 2J^i_{(22)l} - E^i_{(22)l} + 8Kh_{0l}y^i &= -8W^i{}_{||0||0}W_l + 8W^i{}_{||r}W^r{}_{||0}h_{0l} + 8Kh_{0l}y^i,\\
16{}^hR_0{}^i{}_{0l} + 8B^i_{(22)l} - 4D^i_{(22)l} + 2J^i_{(23)l} - E^i_{(23)l} &= 16{}^hR_0{}^i{}_{0l},\\
2J^i_{(32)l} - 4D^i_{(32)l} - E^i_{(32)l} &= 16W^i{}_{||0||0}h_{0l}.
\end{aligned}
$$

Therefore, from (2) of Lemma 3 and the above equalities we have

$$
\begin{aligned}
(3.15)\quad -\frac{1}{4}P^i_{(5)l} &= (h_{00})^2W_0(W^i{}_{||r}W^r{}_{||l} + K\delta^i{}_l) + (h_{00})^2(W^i{}_{||r}W^r{}_{||0}W_l + KW_ly^i)\\
&+2h_{00}(W_0)^2(2W^i{}_{||0||l} - W^i{}_{||l||0}) + 2h_{00}W_0(W^i{}_{||0||0}W_l - W^i{}_{||r}W^r{}_{||0}h_{0l} - Kh_{0l}y^i)\\
&-4(W_0)^3\cdot{}^hR_0{}^i{}_{0l} - 4(W_0)^2W^i{}_{||0||0}h_{0l} = 0,
\end{aligned}
$$

First, we consider the term $(h_{00})^2(W^i{}_{||r}W^r{}_{||0}W_l + KW_ly^i)$ which does not contain W_0. Taking into account that $(h_{00})^2$ is not divisible by W_0, the following equality must hold good

$$W^i{}_{||r}W^r{}_{||0}W_l + KW_ly^i = W_0c_l{}^i(x), \tag{3.16}$$

where $c_l{}^i(x)$ are functions of (x^i) alone, must hold. Transvecting (3.16) by y^l and dividing it by W_0, we get

$$W^i{}_{||r}W^r{}_{||0} + Ky^i = c_l{}^i(x)y^l.$$

Derivating it by y^l, we have

$$W^i{}_{||r}W^r{}_{||l} + K\delta^i{}_l = c_l{}^i(x).$$

Substituting the above equality in (3.16), we get

$$W^i{}_{||r}W^r{}_{||0}W_l + KW_ly^i = W_0(W^i{}_{||r}W^r{}_{||l} + K\delta^i{}_l).$$

Transvecting the above equality by W^l, we get

$$W^i{}_{||r}W^r{}_{||0} + Ky^i = KW_0W^i,$$

where we have used $W^i{}_{||r}W^r{}_{||l}W^l = 0$. Derivating the above equality by y^l, we get

$$W^i{}_{||r}W^r{}_{||l} = KW_lW^i - K\delta^i{}_l. \tag{3.17}$$

Substituting (3.17) in (3.15) and dividing it by $2W_0$, it follows

$$\begin{aligned} &K(h_{00})^2W_lW^i + h_{00}W_0(2W^i{}_{||0||l} - W^i{}_{||l||0}) \\ &\qquad + h_{00}(W^i{}_{||0||0}W_l - KW_0W^ih_{0l}) - 2(W_0)^2 \cdot {}^hR_0{}^i{}_{0l} - 2W_0W^i{}_{||0||0}h_{0l} = 0. \end{aligned} \tag{3.18}$$

Transvecting the above equality by h_{0i}, we get

$$W_{0||l||0} = K(h_{00}W_l - W_0h_{0l}). \tag{3.19}$$

Using $W_{i||j} + W_{j||i} = 0$ and (3.19), we have

$$W_{l||0||0} = -K(h_{00}W_l - W_0h_{0l}). \tag{3.20}$$

Derivating (3.19) by y^i, we have

$$W_{i||l||0} + W_{0||l||i} = K(2h_{0i}W_l - W_ih_{0l} - W_0h_{il}).$$

From the above equality, we have

$$W_{i||0||l} = -W_{i||l||0} - K(2h_{0l}W_i - W_lh_{0i} - W_0h_{li}). \tag{3.21}$$

Using (3.20) and (3.21), the equality (3.18) reduces to

$$3h_{00}(KW_ly^i - Kh_{0l}W^i - W^i{}_{||l||0}) - 2W_0({}^hR_0{}^i{}_{0l} + Ky^ih_{0l} - Kh_{00}\delta^i{}_l) = 0.$$

Since h_{00} is not divisible by W_0, it follows that the equality

$${}^hR_0{}^i{}_{0l} + Kh_{0l}y^i - Kh_{00}\delta^i{}_l = h_{00}d^i{}_l(x), \tag{3.22}$$

where ${d^i}_l(x)$ are functions of (x^i) alone, must hold. Transvecting the above equality by y^l, we get ${d^i}_l(x)y^l = 0$. Derivating the above equality by y^l, we get ${d^i}_l(x) = 0$. Substituting it in (3.22), we get

$$ {}^h{R_0}^i{}_{0l} + Kh_{0l}y^i - Kh_{00}{\delta^i}_l = 0. \tag{3.23}$$

We can rewrite the above equality as

$$ {}^h{R_0}^i{}_{0l} \;=\; K(h_{00}{\delta^i}_l - h_{0l}y^i). $$

Derivating (3.23) by y^j and y^k, we get

$$ {}^h{R_j}^i{}_{kl} + {}^h{R_k}^i{}_{jl} = K(2h_{jk}{\delta^i}_l - h_{jl}{\delta^i}_k - h_{kl}{\delta^i}_j), \tag{3.24}$$

and interchanging j and l, we obtain

$$ {}^h{R_l}^i{}_{kj} + {}^h{R_k}^i{}_{lj} = K(2h_{lk}{\delta^i}_j - h_{lj}{\delta^i}_k - h_{kj}{\delta^i}_l). \tag{3.25}$$

Subtracting (3.25) from (3.24), we get

$$ {}^h{R_j}^i{}_{kl} + 2{}^h{R_k}^i{}_{jl} - {}^h{R_l}^i{}_{kj} = 3K(h_{jk}{\delta^i}_l - h_{kl}{\delta^i}_j). $$

Since the left-hand side of the above equality can be changed as follows:

$$ {}^h{R_j}^i{}_{kl} + 2{}^h{R_k}^i{}_{jl} - {}^h{R_l}^i{}_{kj} = 2{}^h{R_k}^i{}_{jl} - {}^h{R_j}^i{}_{lk} - {}^h{R_l}^i{}_{kj} = 2{}^h{R_k}^i{}_{jl} + {}^h{R_k}^i{}_{jl} = 3{}^h{R_k}^i{}_{jl}, $$

we obtain

$$ {}^h{R_k}^i{}_{jl} = K(h_{jk}{\delta^i}_l - h_{kl}{\delta^i}_j). $$

This means that the Riemannian space (M, h) is of constant curvature K.

Therefore, we obtain

Theorem 2 *Let M be an $n(\geq 2)$-dimensional Riemannian manifold. Put $\alpha = \sqrt{a_{ij}(x)y^iy^j}$ and $\beta = b_i(x)y^i$. Let $(M, \alpha^2/\beta)$ be a Kropina space and define a new Riemannian metric $h = \sqrt{h_{ij}(x)y^iy^j}$ and a vector field W with $|W| = 1$ by (1.2) and (1.3).*

If the Kropina space $(M, \alpha^2/\beta)$ is of constant curvature K, then the vector field W is a Killing one and the Riemannian space (M, h) is of constant curvature K .

3.6 The converse of Theorem 2.

Let $(M, \alpha^2/\beta)$ be a Kropina space and define a new Riemannian metric $h = \sqrt{h_{ij}(x)y^iy^j}$ and a vector field W with $|W| = 1$ by (1.2) and (1.3). Suppose that the vector field W is a Killing one and that the Riemannian space (M, h) is of constant curvature K. To prove that the Kropina space $(M, \alpha^2/\beta)$ is of constant curvature K, we have only to show that the equality (3.9) holds.

Since the vector field W is a Killing one, we have $\mathtt{R}_{00} = 0$. Taking into account (2) of Lemma 2 and (1) of Lemma 3, the Killing part R and the vanishing part Q vanishes respectively.

Therefore, we have only to show that the curvature part $P^i_{(5)l}$ vanishes. At the rest of this subsection, we will prove it. The curvature part $P^i_{(5)l}$ is defined by (3.15).

First, we give the following Lemma 4:

Lemma 4 *For a Killing vector field $W = W^i(\partial/\partial x^i)$ of constant length $|W| = 1$, the equality*

$$W_{i||j||k} = W_r\, {}^hR_k{}^r{}_{ij} \tag{3.26}$$

holds good.

(*Proof.*) From the Ricci's formula, it follows

$$W_{i||j||k} - W_{i||k||j} = -W_r\, {}^hR_i{}^r{}_{jk}. \tag{3.27}$$

On the other hand, since W is a Killing vector field, we have

$$W_{i||j} + W_{j||i} = 0.$$

Applying the h-covariant derivative ${}_{||k}$ to the above equality, we get

$$W_{i||j||k} + W_{j||i||k} = 0.$$

Replacing i, j, k by j, k, i and k, i, j respectively, we obtain the following equalities:

$$\begin{array}{rcl} W_{j||k||i} + W_{k||j||i} &=& 0, \\ W_{k||i||j} + W_{i||k||j} &=& 0. \end{array}$$

Subtracting the second equality from the first equality and adding the third equality to it, we get

$$W_{i||j||k} + W_{i||k||j} - W_r\, {}^hR_j{}^r{}_{ik} - W_r\, {}^hR_k{}^r{}_{ij} = 0. \tag{3.28}$$

From (3.27) and (3.28), we get

$$2W_{i||j||k} = -W_r\, {}^hR_i{}^r{}_{jk} + W_r\, {}^hR_j{}^r{}_{ik} + W_r\, {}^hR_k{}^r{}_{ij}.$$

Using the formula

$${}^hR_i{}^r{}_{jk} + {}^hR_j{}^r{}_{ki} + {}^hR_k{}^r{}_{ij} = 0,$$

we have

$$\begin{array}{rcl} 2W_{i||j||k} &=& W_r\, {}^hR_j{}^r{}_{ki} + W_r\, {}^hR_k{}^r{}_{ij} + W_r\, {}^hR_j{}^r{}_{ik} + W_r\, {}^hR_k{}^r{}_{ij} \\ &=& 2W_r\, {}^hR_k{}^r{}_{ij} \end{array}$$

that is, (3.26) holds good. Q.E.D.

From the assumption that the Riemannian space (M, h) is of constant curvature, we have

$${}^hR_k{}^r{}_{ji} = K(h_{kj}\delta^r{}_i - h_{ki}\delta^r{}_j). \tag{3.29}$$

Using the above equality and (3.26), we get

$$W^i{}_{||j||k} = K(\delta^i{}_k W_j - h_{kj} W^i) \tag{3.30}$$

and from here and ${y^i}_{||j} = 0$ (See, Remark 1), it follows

$$\begin{aligned} {W^i}_{||0||l} &= K({\delta^i}_l W_0 - h_{l0}W^i), \\ {W^i}_{||0||0} &= K(y^i W_0 - h_{00}W^i), \\ {W^i}_{||l||0} &= K(y^i W_l - h_{0l}W^i). \end{aligned} \tag{3.31}$$

From (3.29), we have

$$ {}^h{R_0}^i{}_{0l} = K(h_{00}{\delta^i}_l - h_{0l}y^i) \tag{3.32}$$

and applying the h-covariant derivative $_{||i}$ to the equality $|W|^2 = W_r W_s h^{rs} = 1$, we get

$$W_{r||i}W^r = -W_{i||r}W^r = 0.$$

Furthermore, applying the h-covariant derivative $_{||l}$ to the above equality, we obtain

$$W_{i||r}{W^r}_{||l} + W_{i||r||l}W^r = 0.$$

From the above equality and (3.30), we have

$$\begin{aligned} W_{i||r}{W^r}_{||l} &= -W_{i||r||l}W^r \\ &= K(h_{lr}W_i - h_{li}W_r)W^r \\ &= K(W_l W_i - h_{li}). \end{aligned} \tag{3.33}$$

Substituting the equalities (3.31)-(3.33) in (3.15), we can easily recognize the curvature part $P^i_{(5)\,l} = 0$. Therefore, (3.9) holds good. Hence, from Proposition 2, we get

Theorem 3 *Let M be an $n(\geq 2)$-dimensional Riemannian space. Put $\alpha = \sqrt{a_{ij}(x)y^iy^j}$ and $\beta = b_i(x)y^i$. Let $(M, \alpha^2/\beta)$ be a Kropina space and define a new Riemannian metric $h = \sqrt{h_{ij}(x)y^iy^j}$ and a vector field $W = W^i(\partial/\partial x^i)$ of constant length $|W| = 1$ by (1.2) and (1.3).*

If the vector field $W = W^i(\partial/\partial x^i)$ is a Killing one and the Riemannian space (M, h) is of constant curvature K, the Kropina space $(M, \alpha^2/\beta)$ is of constant curvature K.

From Theorem 2 and Theorem 3, we have

Theorem 4 *Let $(M, \alpha^2/\beta)$ be an $n(\geq 2)$-dimensional Kropina space, where $\alpha^2 = a_{ij}(x)y^iy^j$, $\beta = b_i(x)y^i$ and the matrix (a_{ij}) is positive definite. For the Kropina space, we define a new Riemannian metric $h = \sqrt{h_{ij}(x)y^iy^j}$ and a vector field $W = W^i(\partial/\partial x^i)$ of constant length $|W| = 1$ on M by (1.2) and (1.3).*

Then, the Kropina space $(M, \alpha^2/\beta)$ is of constant curvature K if and only if the following conditions hold:

(1)$W_{i||j} + W_{j||i} = 0$, that is, $W = W^i(\partial/\partial x^i)$ is a Killing vector field.

(2)The Riemannian space (M, h) is of constant curvature K.

Let $(M, F = \alpha^2/\beta)$ be an $n(\geq 2)$-dimensional Kropina space. From Theorem 1, for this Kropina metric $F = \alpha^2/\beta$, we can define a Riemannian metric $h = \sqrt{h_{ij}(x)y^iy^j}$ and a vector field $W = W^i(\partial/\partial x^i)$ of constant length 1 on M by (1.2) and (1.3). We suppose that the vector field W is a Killing one. Then, we have $\mathtt{R}_{00} = 0$. From this assumption, we get the second equation of (3.14), that is, $\mathtt{S}_0 = 0$. Substituting $\mathtt{R}_{00} = 0$, $\mathtt{S}_0 = 0$ and $F = h_{00}/(2W_0)$ in (2.6), we obtain the equation $\Phi^i = -F{\mathtt{S}^i}_0$. Substituing this in (2.4), we get

Theorem 5 *Let $(M, F = \alpha^2/\beta)$ be an $n(\geq 2)$-dimensional Kropina space. We define a Riemannian metric $h = \sqrt{h_{ij}(x)y^i y^j}$ and a vector field $W = W^i(\partial/\partial x^i)$ of constant length $|W| = 1$ on (M, h) by (1.2) and (1.3).*

Suppose that the vector field W is a Killing one, then the coefficients G^i of the geodesic spray of the Kropina space $(M, \alpha^2/\beta)$ is written as follows:

$$2G^i \;=\; {}^h\gamma_0{}^i{}_0 - 2F\boldsymbol{S}^i{}_0, \tag{3.34}$$

where ${}^h\gamma_j{}^i{}_k$ are Christoffel symbols of the Riemannian space (M, h).

Remark 2 *The geodesic spray of the Randers space $(M, \alpha + \beta)$ is given in the subsection 2.3 ([5] p.5). Comparing to this, the geodesic spray of the Kropina space $(M, F = \alpha^2/\beta)$ is in a very simple form (3.33).*

Acknowledgments. The authors would like to express their sincere thanks to Professor Dr. S. V. Sabau and Professor Dr. H. Shimada for the valuable advices, the efforts to check all the calculation and the continuous encouragements.

References

[**1**] S. Bácsó, X. Cheng and Z. Shen : Curvature properties of (α, β)-metrics, Advanced Studies in Pure Mathematics 48, 2007, Finsler Geometry, Sappro 2005 - In Memory of Makoto Matsumoto, 73-110.

[**2**] M. Matsumoto : Foundations of Finsler geometry and special Finsler spaces, Kaiseisha Press, Saikawa, Otsu, Japan, 1986.

[**3**] M. Matsumoto: Finsler spaces of constant curvature with Kropina metric, Tensor N.S., 50(1991), 194-201.

[**4**] M. Matsumoto: Theory of Finsler spaces with $(\alpha, \beta)-$metric, Rep. Math. Phys., 31(1992), 43-83.

[**5**] C. Robles : Geodesic in Randers spaces of constant curvature, arXiv:math/0501358v2 [math.DG] 8 July 2005.

[**6**] S. Tachibana : Riemannian geometry (Japanese), Modern mathematical lecture 15, Asakura Shoten, Tokyo, 1967.

[**7**] R. Yoshikawa and K. Okubo: Kropina spaces of constant curvature, Tensor N.S., 68(2007), 190-203.

Hachiman technical
High School
5 Nishinosho-cho Hachiman
523-0816 Japan
E-mail: ryozo@e-omi.ne.jp

Faculty of Education
Shiga University
2-5-1 Hiratsu Otsu
520-0862 Japan
E-mail: okubo@edu.shiga-u.ac.jp